

Proceedings of the Iowa Academy of Science

Volume 32 | Annual Issue

Article 34

1925

Some Wound Responses of Citrus Leaves

Robert B. Wylie
State University of Iowa

Let us know how access to this document benefits you

Copyright ©1925 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Wylie, Robert B. (1925) "Some Wound Responses of Citrus Leaves," *Proceedings of the Iowa Academy of Science*, 32(1), 265-265.

Available at: <https://scholarworks.uni.edu/pias/vol32/iss1/34>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

SOME WOUND RESPONSES OF CITRUS LEAVES

ROBERT B. WYLIE

(*ABSTRACT*)

The general wound response of the foliar organs of Citrus was studied using leaves of a large potted lemon tree. This plant stood at the southeast corner of a building, outdoors, during the experimental period and received full sunshine a large part of the day. Following wounding an interval of about ten days elapsed before there was a division of the living cells beneath the collapsed dead tissue. The cicatrice, which was not complete until the close of the third week following wounding, consisted of eight to ten layers of cells.

STATE UNIVERSITY OF IOWA.

SOME NOTEWORTHY IOWA FUNGI OF 1925

GUY WEST WILSON

(*ABSTRACT*)

Cercospora on potato, *Peronospora Viciae* on garden peas, *Colosporium Solidaginis* on cultivated aster, and the aecidial stages of *Puccinia Sorghi* on *Oxalis stricta* were noted during the season in the vicinity of Fayette, Iowa.

UPPER IOWA UNIVERSITY.

MICROSPOROGENESIS IN CUCURBITA MAXIMA

EDWARD F. CASTETTER

(*ABSTRACT*)

A pure line of Hubbard squash was used for the investigation. In the development of the anther the pollen mother cells are formed from a single longitudinal row of cells of the primary sporogenous layer in each anther lobe. During early diakinesis a perinuclear zone is formed. From the perinuclear zone as well as from the interior of the nucleus fibers are formed which con-